

Original Article

Molecular Interactions Study in The Solution of Root Extract of *Ricinus Communis* At 3 MHz

 Rahul N. Gaikwad¹, Shailesh S. Kamble²
¹Department of chemistry, Shree V. R. College, Sawana, Dist. Yavatmal, Maharashtra

²Department of Chemistry, Arts, commerce and Science College, Maregaon, Maharashtra

Manuscript ID:

IBMIIRJ -2026-030512

Submitted: 10 Apr. 2026

Revised: 20 Apr. 2026

Accepted: 08 May 2026

Published: 31 May 2026

ISSN: 3065-7857

Volume-3

Issue-5

Pp. 72-75

May 2026

Correspondence Address:

 Rahul N. Gaikwad
 Department of chemistry, Shree V. R.
 College, Sawana, Dist. Yavatmal,
 Maharashtra.
 Email:
raahulgaikwad955276@gmail.com


Quick Response Code:


 Web: <https://ibrj.us>


DOI: 10.5281/zenodo.21696664

DOI Link:

<https://doi.org/10.5281/zenodo.21696664>


Creative Commons

Abstract

Ultrasonic Velocity, density, viscosity have been measured experimentally for the solution of leaf extract of *Ricinus communis* in 50% ethyl alcohol with various concentrations at 298.15 K, 303.15 K, 308.15 K keeping constant frequency of 4 MHz. As the acoustical parameters like adiabatic compressibility, intermolecular free length, relative association, relaxation time, specific acoustic impedance would prove to be more useful to predict and confirm the molecular interactions, these have been determined by measuring the Ultrasonic Velocity, density, viscosity of the prepared solution. A variation in these parameters will provide strong information regarding the molecular interactions taking place in the solution. The propagation of ultrasonic waves affects the physical properties of the medium and hence gives useful information about the molecular interactions such as ion-ion, ion-dipole, dipole-dipole interaction, hydrogen bonding in the solution. Here the study shows that the ultrasonic velocity increases here with increase in concentration this is due to the solvation process of solute by two different solvent molecules like water and ethanol, from this we can say that there is strong solute-solvent interaction in the root extract solution of *Ricinus communis*.

Keywords: - Ultrasonic velocity, Adiabatic compressibility, Relative association, Intermolecular free length.

Introduction

Ultrasonic velocity measurements and other acoustical parameters of liquid mixtures are the powerful technique in understanding the chemical nature and the molecular interactions [1-4]. Many researchers used ultrasonic velocity measurement for studying solute-solvent interaction in number of systems including organic liquid, dilute solutions in organic acid and complexes [5-7]. Ultrasonic velocity in liquids and liquid mixtures provide valuable information about their physico-chemical properties and the nature of molecular interactions in them [8-10]. Our country is very well known for Ayurveda, in the Ayurveda medicines are largely made up from plants, herbs. One of such plants is which is also known as *Ricinus communis*. *Ricinus communis* is known for its antifungal nature. The solution of leaf extract of *Ricinus communis* in 50% ethyl alcohol is studied at 4 MHz for the concentration of 1%, 0.5%, 0.25%, 0.125% at 298.15K, 303.15K, 308.15K. Here the effect of concentration at different temperature on molecular interaction will be predicted which may be helpful for predicting the reactivity of the extract.

Materials And Methods

The leaf extract used in this study was of analytical range. 50% ethyl alcohol was used for the preparation of solution. A special thermostatic water bath arrangement was made to maintain constant temperature. 1%, 0.5%, 0.25%, 0.125% solutions of leaf extracts of *Ricinus communis* was Prepared by taking accurate weights on electronic digital balance (Model CB/CA/CT-Series, Contech having accuracy ± 0.0001 g.) The ultrasonic velocity of the 1%, 0.5%, 0.25%, 0.125% solutions of leaf extracts of *Ricinus communis* was measured with the Multifrequency ultrasonic interferometer (Model M-83, Mittal Enterprises) at 3 MHz frequency with an accuracy of ± 2 m/s. All the readings were taken at 298.15 K, 303.15K, 308.15K.

Creative Commons (CC BY-NC-SA 4.0)

This is an open access journal, and articles are distributed under the terms of the [Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International](https://creativecommons.org/licenses/by-nc-sa/4.0/) Public License, which allows others to remix, tweak, and build upon the work noncommercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

How to cite this article:

Gaikwad1, R. N. & Kamble, S. S. (2026). Molecular Interactions Study in The Solution of Root Extract of *Ricinus Communis* At 3 MHz. *Insight Bulletin: A Multidisciplinary Interlink International Research Journal*, 3(5), 72–75. <https://doi.org/10.5281/zenodo.21696664>

The viscosity was measured by using Ostwald's viscometer and the density of the solution was measured by using Digital densitometer (DMA-35, Anton paar).

Computation

By using ultrasonic velocity following ultrasonic parameters are calculated.

1. Adiabatic compressibility -

$$\beta = 1/v_s^2 d$$

Where, v - velocity of solution,

d - density of liquid

2. Intermolecular free length -

$$L_r = K\sqrt{\beta}$$

Where, K - temperature dependent known as Jacobson's constant

3. Specific acoustic impedance -

$$Z = v \times d_s$$

4. Relative association -

$$R_a = d_s / d_o [v_o/v_s]^{1/3}$$

Where, v_o - ultrasonic velocity of solvent

v_s - ultrasonic velocity of solution

5. Relaxation time -

$$\tau = 4/3 \beta \times \eta$$

Result And Discussion







The experimentally determined values are listed in the following table.

Table 1: Density, Viscosity and Velocity (at frequency 3 MHz) of *Ricinus communis* leaf extract solution.

SR.NO.	Conc.	Temp. (K)	Density (d_s)	Velocity (v_s) (m/s)	Viscosity (η)
	(%)		(Kg m ⁻³)		(Kg m ⁻¹ s ⁻²)
1)	1%	298.15	934.1	1616.8	19.78 E ⁻⁴
		303.15	931	1504.7	20.651 E ⁻⁴
		308.15	924.3	1501.2	16.26 E ⁻⁴
2)	0.50%	298.15	933	1540.3	20.42 E ⁻⁴
		303.15	929.3	1583	19.79 E ⁻⁴
		308.15	925.2	1681.2	17.07 E ⁻⁴
3)	0.25%	298.15	929.3	1502.1	16.73 E ⁻⁴
		303.15	926	1531.2	17.66 E ⁻⁴
		308.15	923	1608.1	15.207 E ⁻⁴
4)	0.13%	298.15	928.2	1500.7	16.41 E ⁻⁴
		303.15	925.6	1515.2	17.28 E ⁻⁴
		308.15	922.2	1601.1	15.00 E ⁻⁴

Table 2: - Acoustic parameters of *Ricinus Communis* leaf extract solution in 50% Ethyl Alcohol at 3 MHz.

Sr. No.	Conc. (%)	Temp. (K)	Adiabatic Compressibility	Specific Acoustic Impedence Kg M ⁻² S ⁻¹	Intermolecular free length	Relative Association	Relaxation time
1.	1%	298.15	4.095 E ⁻¹⁰	1510252.8	4.162 E ⁻¹¹	3.600 E ⁻¹	10.799 E ⁻¹³
		303.15	4.744 E ⁻¹⁰	1400875.7	4.520 E ⁻¹¹	4.256 E ⁻¹	13.029 E ⁻¹³
		308.15	4.8007 E ⁻¹⁰	1387559.1	4.5887 E ⁻¹¹	5.265 E ⁻¹	10.214 E ⁻¹³
2.	0.5%	298.15	4.517 E ⁻¹⁰	1437099	4.371 E ⁻¹¹	3.775 E ⁻¹	12.26 E ⁻¹³
		303.15	4.294 E ⁻¹⁰	1471081	4.301 E ⁻¹¹	4.041 E ⁻¹	11.30 E ⁻¹³
		308.15	3.884 E ⁻¹⁰	1543326	4.127 E ⁻¹¹	4.742 E ⁻¹	8.81 E ⁻¹³
3.	0.25%	298.15	4.769 E ⁻¹⁰	1395901.5	4.491 E ⁻¹¹	3.856 E ⁻¹	10.61 E ⁻¹³
		303.15	4.606 E ⁻¹⁰	1417706	4.454 E ⁻¹¹	4.162 E ⁻¹	10.58 E ⁻¹³
		308.15	4.189 E ⁻¹⁰	1484276	4.286 E ⁻¹¹	4.907 E ⁻¹	8.35 E ⁻¹³
4.	0.125 %	298.15	4.795 E ⁻¹⁰	1389648.2	4.503 E ⁻¹¹	3.846 E ⁻¹	10.46 E ⁻¹³
		303.15	4.713 E ⁻¹⁰	1400044.8	4.500 E ⁻¹¹	4.194 E ⁻¹	10.83 E ⁻¹³
		308.15	4.24 E ⁻¹⁰	1472920	4.312 E ⁻¹¹	4.913 E ⁻¹	8.45 E ⁻¹³



The ultrasonic velocity of 1%, 0.5%, 0.25%, 0.125% Ricinus Communis (Erandi) root extract was measured at 298.15K, 303.15K, 308.15K at 4 MHz frequency. From Table no.1 & fig.no.1, shows that ultrasonic velocity increases with increase in concentration this is because of the solvation process of solute by two different solvent molecules that is water and ethyl alcohol, it is observed that the ultrasonic velocity increases with the increase in temperature this is because of structure in liquid mixture with many cavities. These cavities can accommodate solute molecules. The ultrasonic velocity increases here with increase in concentration this is due to the solvation process of solute by two different solvent molecules like water and ethanol. From table no. 2 and fig.2 it is observed that the adiabatic compressibility decreases with increase in concentration this may be attributed to increase in compressible molecules. A decreased value of adiabatic compressibility indicates strong interaction between components of the mixture. From Fig.3 it is seen that the specific acoustic impedance also increases with increase in concentration, this increase in acoustic impedance shows increase in molecular packing in the medium which gives the possibility of molecular interaction due to hydrogen bonding between solute and solvent. Fig.4. shows that intermolecular free length increases with decrease in concentration this is due to closed packed structure formed by one solute and two solvent molecules.

Fig.5. shows decrease in relative association with increase in concentration this is mainly due to breaking of associated solvent-solvent molecules on addition of solute in it. Fig 6. shows that as concentration increases relaxation time increases it is because of the relaxation process showing the presence of strong molecular interactions.

Conclusion

The ultrasonic velocity increases here with increase in concentration this is due to the solvation process of solute by two different solvent molecules like water and ethanol. Evaluation of adiabatic compressibility, specific acoustic impedance, relative association, relaxation time, intermolecular free length clearly shows strong solute solvent interaction.

Acknowledgment

The authors express their sincere gratitude to their respective institutions for providing the necessary facilities and support to carry out this research work.

Financial support and sponsorship

Nil.

Conflicts of interest

The authors declare that there are no conflicts of interest regarding the publication of this paper.

References

1. Praveen, S., Shukla, D., Singh, S., Singh, K. P., & Shukla, J. P. (2009). *App Acous.*, **70**(Suppl 3), 507.
2. Umadevi, M., Keshavswamy, R., Rathina, K., & Mahalaxmi, R. (2016). *Journal of Molecular Liquids*, **219**, 820.
3. Nayak, N., Aralaguppi, M. I., & Aminabhavi, T. M. (2003). *Journal of Chemical & Engineering Data*, **48**, 1489–1494.
4. Steele, W. V., Chirico, R. D., Cowell, A. B., Knipmeyer, S. E., & Nguyen, A. (2002). *Journal of Chemical & Engineering Data*, **47**(Suppl 4), 667–688.
5. Reis, J. C. R., Santos, A. F. S., Disas, F. A., & Lampreia, I. M. S. (2008). *ChemPhysChem*, **9**, 1178–1188.
6. Khansanshin, T. S., Shchemelev, A. P., Srivastava, T. N., Singh, R. P., & Swaroop, B. (1983). *Indian Journal of Pure Physics*, **21**, 67–72.
7. Kumar, A., Srivastava, U., Singh, A. K., & Shukla, R. K. (2016). 4(Suppl 2), 157–167. (*Journal name missing—please confirm*)
8. Bhatt, S. C., Semwak, H., Lingwal, V., Singh, K., & Semwak, B. S. (2000). *Journal of the Acoustical Society of India*, **28**, 293–296.
9. Dhanlashmi, A., & Sekar, S. (1999). *Journal of Pure and Applied Ultrasonics*, **21**, 97.
10. Sastry, G. L. N. (1986). *Journal of the Acoustical Society of India*, **14**, 1–9.